

Similac® 360 Total Care® Sensitive Liquid

Milk-based infant formula with iron

SIMILAC 360 TOTAL CARE SENSITIVE is a 20 Cal/fl oz, nutritionally complete, non-GMO*, milk-based, iron fortified infant formula designed for sensitive tummies as a milk-based alternative to standard milk-based formulas. Suitable for infants with lactose sensitivity.

* Ingredients not genetically engineered.



Features

- Designed to support the immune system, brain development, and digestive health.
- HMOs are immune nourishing prebiotics structurally identical to those in breast milk.
- DHA, lutein, vitamin E.
- Similac 360 Total Care Sensitive for fussiness and gas due to lactose sensitivity.
- No artificial growth hormones. †
- Non-GMO. *
- No palm olein oil.
- Halal.
- Kosher; made with dairy ingredients.

* Ingredients not genetically engineered.

† Milk ingredients are derived from milk of non-rbST-treated cows, which is not different from milk of rbST-treated cows.

Safety Precautions

- Not for infants with galactosemia.
- Never use a microwave oven to warm formula. Serious burns can result.

Availability

List Number	Item Description
68155	Similac 360 Total Care Sensitive Ready to Feed / 2 fl oz (59 mL) Bottle / 48 ct
68154	Similac 360 Total Care Sensitive Ready to Feed / 2 fl oz (59 mL) Bottle / 48 ct
68152	Similac 360 Total Care Sensitive Ready to Feed / 2 fl oz (59 mL) Bottle / 2 x 12 ct
68149	Similac 360 Total Care Sensitive Ready to Feed / 2 fl oz (59 mL) Bottle / 2 x 12 ct

Ingredients

Unflavored Liquid: Water, Maltodextrin, Milk Protein Isolate, High Oleic Safflower Oil, Sugar, Soy Oil, Coconut Oil. Less than 0.5% of: Human Milk Oligosaccharides (2'-Fucosyllactose, Lacto-N-Tetraose, 3-Fucosyllactose, 6'-Sialyllactose, 3'-Sialyllactose)*, Schizochytrium Sp. Oil†, M. Alpina Oil‡, Beta-Carotene, Lutein, Calcium Phosphate, Potassium Citrate, Carrageenan, Ascorbic Acid, Potassium Chloride, Monoglycerides, Soy Lecithin, Magnesium Chloride, Inositol, Magnesium Phosphate, Choline Chloride, Calcium Carbonate, Ferrous Sulfate, Choline Bitartrate, Taurine, Zinc Sulfate, L-Carnitine, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Vitamin B12, Sodium Citrate, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate).* HMO, not from human milk.† Source of DHA.‡ Source of ARA.

Contains milk and soy ingredients.

Nutrition Information

Nutrient	100 Cal (5 fl oz)		1000 mL (1.1 QT)	
	Value	%DV	Value	%DV
Calories	100		676	
Volume, mL	148		1000	
Protein, g	2.1		14.19	
Fat, g	5.4		36.49	
Carbohydrate, g	10.9		73.65	
Water, g	133		899	
Linoleic Acid, mg	1000		6757	
Potential Renal Solute Load, mOsm	19.7		132.8	
Vitamin A, IU	300		2027	
Vitamin D, IU	60		405	
Vitamin E, IU	1.5		10.1	
Vitamin K, mcg	8		54.1	
Thiamin (Vitamin B1), mcg	100		676	
Riboflavin (Vitamin B2), mcg	150		1014	
Vitamin B6, mcg	60		405	
Vitamin B12, mcg	0.25		1.69	
Niacin, mcg	1050		7095	

Similac® 360 Total Care® Sensitive Liquid

Nutrient	100 Cal (5 fl oz)		1000 mL (1.1 QT)	
	Value	%DV	Value	%DV
Folic Acid (Folacin), mcg	15		101.4	
Pantothenic Acid, mcg	450		3041	
Biotin, mcg	4.4		29.7	
Vitamin C (Ascorbic Acid), mg	9		60.8	
Choline, mg	24		162.2	
Inositol, mg	24		162.2	
Calcium, mg	84		568	
Calcium, mEq	4.2		28.4	
Phosphorus, mg	56		378	
Magnesium, mg	6		40.5	
Iron, mg	1.8		12.16	
Zinc, mg	0.75		5.07	
Manganese, mcg	5		33.8	
Copper, mcg	90		608	
Iodine, mcg	15		101	
Selenium, mcg	2		14	
Sodium, mg	30		203	
Sodium, mEq	1.3		8.8	
Potassium, mg	107		723	
Potassium, mEq	2.7		18.2	
Chloride, mg	65		439	
Chloride, mEq	1.8		12.2	



Preparation

Instructions for Use:

Your baby's health depends on carefully following these directions. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to boil (sterilize) nipples and rings before use. Ready to feed formula if mixed with water may not provide proper nutrition, and repeated use of such feedings could cause illness.

- Do not add water.Shake very well.
- Do not add water.
- Shake very well.

2 or 8 fl oz bottle

- Do not use if breakaway ring is missing or broken.
Twist off cap.Use clean hands to twist on clean nipple and ring. Use sterile nipple and ring if directed.No need to warm before feeding.
For use in a different feeding container (cup or bottle):
Ask your baby's doctor if you need to boil (sterilize) the empty bottles before adding formula.Pour desired amount into clean feeding container. Recap bottle, refrigerate and use within 48 hours.
Once feeding begins, use within 1 hour or discard.
- Do not use if breakaway ring is missing or broken.
Twist off cap.Use clean hands to twist on clean nipple and ring. Use sterile nipple and ring if directed.No need to warm before feeding.
- Twist off cap.
- Use clean hands to twist on clean nipple and ring. Use sterile nipple and ring if directed.
- No need to warm before feeding.
- For use in a different feeding container (cup or bottle):
Ask your baby's doctor if you need to boil (sterilize) the empty bottles before adding formula.Pour desired amount into clean feeding container. Recap bottle, refrigerate and use within 48 hours.
- Ask your baby's doctor if you need to boil (sterilize) the empty bottles before adding formula.
- Pour desired amount into clean feeding container. Recap bottle, refrigerate and use within 48 hours.
- Once feeding begins, use within 1 hour or discard.

32 fl oz bottle

- Do not use if band around cap or inner foil seal is damaged.Shake very well before each use.Remove protective band; twist off and clean cap.Invert cap; press down to pierce foil, then turn cap a half turn.Remove foil.Pour formula into bottle; attach clean nipple.Once feeding begins, use within 1 hour or discard.Prepare and feed all formula in the container within 3 days after opening.
- Do not use if band around cap or inner foil seal is damaged.

- Shake very well before each use.
- Remove protective band; twist off and clean cap.
- Invert cap; press down to pierce foil, then turn cap a half turn.
- Remove foil.
- Pour formula into bottle; attach clean nipple.
- Once feeding begins, use within 1 hour or discard.
- Prepare and feed all formula in the container within 3 days after opening.

STORAGE INSTRUCTIONS:

2 & 8 fl oz Bottle

- Store unopened at room temperature; avoid extreme temperatures. Do not reuse bottle. Avoid prolonged exposure of bottles to light. Store in carton until ready for use.
- Store unopened at room temperature; avoid extreme temperatures.
- Do not reuse bottle.
- Avoid prolonged exposure of bottles to light.
- Store in carton until ready for use.

32 fl oz Bottle

- Unopened Container: Store at room Temperature; avoid extreme Temperatures. Opened Container: Recap, refrigerate, and feed within 3 days. Prepared unfed bottles: cap, refrigerate, and feed within 2 days. Do not reuse container.
- Unopened Container: Store at room Temperature; avoid extreme Temperatures.
- Opened Container: Recap, refrigerate, and feed within 3 days.
- Prepared unfed bottles: cap, refrigerate, and feed within 2 days. Do not reuse container.