

Similac® Advance®

Infant formula with iron

SIMILAC ADVANCE is a 20 Cal/fl oz, nutritionally complete, milk-based, iron-fortified infant formula for use as an alternative or supplement to breastfeeding.



Features

- OptiGRO® is our exclusive blend of DHA, lutein and vitamin E: these important ingredients are found in breast milk.
 - DHA for brain and eye development.
 - Lutein to support eye health.
 - Vitamin E to support developing cells.
- Our exclusive formula has:
 - Calcium for strong bones - no palm olein oil.
 - Nucleotides to help support the immune system.
 - Prebiotics to help promote digestive health.
 - Carotenoids like those naturally found in breast milk.
- Clinically proven tolerance in newborns. 1
- Clinically shown to support greater calcium absorption * for strong bones. 2,3
- Gluten-free.
- Kosher; made with dairy ingredients.
- Halal.

*• Similac® with Iron vs Enfamil® with Iron. Enfamil is not a trademark of Abbott Laboratories.

1• Lloyd B, et al. Pediatrics. 1999;103(1):1-6.

2• Nelson SE, et al. J Am Coll Nutr. 1998;17(4):327-332.

3• Koo W, et al. Pediatrics. 2003;111:1017-1023.

Safety Precautions

- Not for infants or children with galactosemia.
- Never use a microwave oven to warm formula. Serious burns can result.
- Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.

Availability

List Number

53363

Item Description

Similac Advance Ready to Feed / 32 fl oz (946 mL) Bottle / 6 ct

Ingredients

Unflavored Liquid: Water, Nonfat Milk, Lactose, High Oleic Safflower Oil, Soy Oil, Coconut Oil, Galactooligosaccharides*, Whey Protein Concentrate. Less than 0.5% of: Schizochytrium Sp. Oil†, M. Alpina Oil‡, Beta-Carotene, Lutein, Ascorbic Acid, Calcium Carbonate, Potassium Citrate, Soy Lecithin, Monoglycerides, Potassium Chloride, Carrageenan, Magnesium Chloride, Ferrous Sulfate, Choline Bitartrate, Choline Chloride, Taurine, Calcium Phosphate, Potassium Phosphate, Inositol, Zinc Sulfate, Niacinamide, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, L-Carnitine, Riboflavin, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Vitamin B12, Salt, Potassium Iodide, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate). * Prebiotic.† Source of DHA.‡ Source of ARA.

Contains milk and soy ingredients.

Nutrition Information

Nutrient	100 Cal (5 fl oz)		1000 mL (1.1 QT)	
	Value	%DV	Value	%DV
Calories	100		676	
Volume, mL	148		1000	
Protein, g	2.07		14	
Fat, g	5.4		36.5	
Carbohydrate, g	11.2		75.7	
Water, g	134		905	
Linoleic Acid, mg	1000		6757	
Potential Renal Solute Load, mOsm	18.7		126.5	
Vitamin A, IU	300		2027	
Vitamin D, IU	60		405	
Vitamin E, IU	1.5		10.1	
Vitamin K, mcg	8		54.1	

Similac® Advance®

Nutrient	100 Cal (5 fl oz)		1000 mL (1.1 QT)	
	Value	%DV	Value	%DV
Thiamin (Vitamin B1), mcg	100		676	
Riboflavin (Vitamin B2), mcg	150		1014	
Vitamin B6, mcg	60		405	
Vitamin B12, mcg	0.25		1.69	
Niacin, mcg	1050		7095	
Folic Acid (Folacin), mcg	15		101.4	
Pantothenic Acid, mcg	450		3041	
Biotin, mcg	4.4		29.7	
Vitamin C (Ascorbic Acid), mg	9		60.8	
Choline, mg	24		162.2	
Inositol, mg	24		162.2	
Calcium, mg	78		527	
Calcium, mEq	3.9		26.4	
Phosphorus, mg	42		284	
Magnesium, mg	6		40.5	
Iron, mg	1.8		12.16	
Zinc, mg	0.75		5.07	
Manganese, mcg	5		33.8	
Copper, mcg	90		608	
Iodine, mcg	15		101	
Selenium, mcg	2		14	
Sodium, mg	24		162	
Sodium, mEq	1		6.8	
Potassium, mg	105		709	
Potassium, mEq	2.7		18.2	
Chloride, mg	65		439	
Chloride, mEq	1.8		12.2	

Preparation

INSTRUCTIONS FOR USE:

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

Powder

1. Wash your hands, surfaces and utensils. Pour water into clean bottle.
2. Add 1 unpacked level scoop (8.7 g) to each 2 fl oz of water. Return dry scoop to container.
3. Cap bottle; shake well; attach clean nipple. Once feeding begins, use within 1 hour or discard.

Powder Mixing Guide

Measure water	+	Add scoop(s) of unpacked level powder using enclosed scoop
2 fl oz		1 scoop (8.7 g)
4 fl oz		2 scoops
6 fl oz		3 scoops
8 fl oz		4 scoops

Each scoop adds about 0.2 fl oz to the amount of prepared formula. For larger size mixing instructions, please visit www.Similac.com/mixing info.

When mixed as directed:

20.6 oz container makes approximately 150 oz of formula.

Ready to Feed

- Do not add water.
 - Do not use if band around cap or inner foil seal is damaged.
1. Shake very well before each use. Remove protective band, twist off and clean cap.
 2. Invert cap; press down to pierce foil, then turn cap a half turn. Remove foil.
 3. Pour formula into bottle; attach clean nipple. Once feeding begins, use within 1 hour or discard.
- Prepare and feed all formula from the 1-QT container within 3 days after opening.

Concentrated Liquid



1. Rinse can lid; shake very well; open with a clean punch-type opener.
2. Pour equal amounts of formula and water into bottle.
3. Attach nipple; swirl gently to mix. Once feeding begins, use within 1 hour or discard.

STORAGE AND HANDLING:

Powder

- Once mixed, store bottles in refrigerator and feed to baby within 24 hours.
- Store unopened or opened container at room temperature; avoid extreme temperatures.
- Use opened container contents within 1 month.

Ready to Feed

- Once opened, store quart bottle in refrigerator.
- Store prepared bottles in refrigerator and feed to baby within 48 hours.
- Store unopened at room temperature; avoid extreme temperatures.
- Do not reuse container.
- Protect from light.

Concentrated Liquid

- Cover opened cans and prepared formula not used immediately. Store in refrigerator and feed to baby within 48 hours.
- Store unopened cans at room temperature; avoid extreme temperatures.